



Australian Government
Department of Health

COMMUNICABLE DISEASES INTELLIGENCE

2019 Volume 43
<https://doi.org/10.33321/cdi.2019.43.24>

Evaluation of enhanced follow-up for pertussis in children aged 5–9 years in Sydney Local Health District, NSW

Kwendy Cavanagh, Emma Quinn, Alexandre S Stephens, Zeina Najjar, Essi Huhtinen
and Leena Gupta

Communicable Diseases Intelligence

ISSN: 2209-6051 Online

This journal is indexed by Index Medicus and Medline.

Creative Commons Licence - Attribution-NonCommercial-NoDerivatives CC BY-NC-ND

© 2019 Commonwealth of Australia as represented by the Department of Health

This publication is licensed under a Creative Commons Attribution-Non-Commercial NoDerivatives 4.0 International Licence from <https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode> (Licence). You must read and understand the Licence before using any material from this publication.

Restrictions

The Licence does not cover, and there is no permission given for, use of any of the following material found in this publication (if any):

- the Commonwealth Coat of Arms (by way of information, the terms under which the Coat of Arms may be used can be found at www.itsanhonour.gov.au);
- any logos (including the Department of Health's logo) and trademarks;
- any photographs and images;
- any signatures; and
- any material belonging to third parties.

Disclaimer

Opinions expressed in Communicable Diseases Intelligence are those of the authors and not necessarily those of the Australian Government Department of Health or the Communicable Diseases Network Australia. Data may be subject to revision.

Enquiries

Enquiries regarding any other use of this publication should be addressed to the Communication Branch, Department of Health, GPO Box 9848, Canberra ACT 2601, or via e-mail to: copyright@health.gov.au

Communicable Diseases Network Australia

Communicable Diseases Intelligence contributes to the work of the Communicable Diseases Network Australia.
<http://www.health.gov.au/cdna>



Communicable Diseases Intelligence (CDI) is a peer-reviewed scientific journal published by the Office of Health Protection, Department of Health. The journal aims to disseminate information on the epidemiology, surveillance, prevention and control of communicable diseases of relevance to Australia.

Editor

Cindy Toms

Deputy Editor

Simon Petrie

Design and Production

Kasra Yousefi

Editorial Advisory Board

David Durrheim,
Mark Ferson, John Kaldor,
Martyn Kirk and Linda Selvey

Website

<http://www.health.gov.au/cdi>

Contacts

Communicable Diseases Intelligence is produced by:
Health Protection Policy Branch
Office of Health Protection
Australian Government
Department of Health
GPO Box 9848, (MDP 6)
CANBERRA ACT 2601

Email:

cdi.editor@health.gov.au

Submit an Article

You are invited to submit your next communicable disease related article to the Communicable Diseases Intelligence (CDI) for consideration. More information regarding CDI can be found at:
<http://health.gov.au/cdi>.

Further enquiries should be directed to:

cdi.editor@health.gov.au.

Short report

Evaluation of enhanced follow-up for pertussis in children aged 5–9 years in Sydney Local Health District, NSW

Kwendy Cavanagh, Emma Quinn, Alexandre S Stephens, Zeina Najjar, Essi Huhtinen and Leena Gupta

Background

Pertussis remains endemic in Australia despite a long-term national vaccination program, with waning immunity¹ contributing to epidemics every 3–5 years.² Pertussis is a notifiable disease in NSW³ and public health units (PHUs) actively follow-up cases in children aged 0–4 years and women in the last month of pregnancy to protect those most at-risk of severe disease, i.e. infants aged less than six months.⁴ This follow-up aims to prevent transmission to infants by ensuring timely treatment of cases and contact tracing to identify any high-risk close contacts who require post-exposure prophylaxis (PEP).⁴

During the 2010–2011 epidemic in NSW, notification rates were highest in children aged 5–9 years in both NSW⁵ and in Sydney Local Health District (SLHD), an inner-Sydney region (Figure 1). In late 2014, pertussis notifications resurged across NSW and SLHD, again predominantly in school-aged children (Figure 1). Evidence demonstrates that both parents^{6,7} and siblings⁸ are potential sources of infant pertussis. More recent data also suggest that school-aged children are most likely to introduce pertussis to households.⁹ Therefore, in response to this evidence, SLHD PHU commenced enhanced surveillance of children aged 5–9 years during 2015.


Keywords: pertussis, source of illness, pertussis vaccination, contacts

Methods

We evaluated the effectiveness of the enhanced follow-up in children aged 5–9 years in SLHD during 2015 by determining: (i) differences in notification and hospitalisation rates in children aged 0–9 years in 2015 compared to 2011; (ii) the key sources of illness for cases aged 0–4 years in 2015 compared to 2011; and (iii) the incremental cost-benefit of following up children aged 5–9 years, by calculating the ratio of additional cases with high-risk contacts identified for PEP per unit of PHU staff time.

Numbers and rates of notifications and hospitalisations related to pertussis for the two epidemic years (2011 vs 2015) in SLHD were compared using the NSW Communicable Disease Register (CDR). These periods were chosen because children aged 0–9 years were offered the same vaccination schedule. The CDR links data from the Notifiable Conditions Incident Management System (NCIMS) with the Admitted Patient Data Collection (APDC), amongst other data. Confirmed pertussis cases in SLHD were identified based on condition code, case classification and geography of residence. Hospitalisations related to pertussis were identified by merging confirmed notifications with APDC data where the hospitalisation date was within 22 days of the notification date.¹⁰ Only hospitalisations with a respiratory-related cause based on

Figure 1. Age-specific rates of pertussis by age group per 100,000 population in Sydney Local Health District (SLHD)



Source: Data from the Notifiable Conditions Incident Management System from within the NSW Communicable Disease Register, accessed through the Secure Analytics for Population Health Research and Intelligence (SAPHaRI) portal. Date of extraction and analysis 10/05/2017.

ICD-10 codes, excluding nested transfers were counted. Age-specific rates were calculated using mid-year population estimates from the Australian Bureau of Statistics. CDR data were analysed using SAS Enterprise Guide software, Version 9.3™.

PHUs routinely follow-up pertussis cases aged 0–4 years by interviewing both the doctor and the caregiver to collect epidemiological data which is entered into NCIMS⁴. During the study period of 1 November 2014 to 1 November 2015, SLHD PHU also followed up cases of pertussis in children aged 5–9 years using the same process. Sources of illness for cases aged 0–4 years were extracted from NCIMS and analysed in Excel. NCIMS data on cases in children aged 0–9 years in 2015 were audited to ascertain the existence of high-risk contacts requiring PEP. The average time to complete one case was estimated by PHU staff. Ethical approval was obtained from the SLHD Human Research Ethics Committee under section 5.1.22 of the National Statement on Ethical Conduct in Human Research.¹¹ The

CDR is made available under the Public Health and Diseases Registers provisions of the NSW Public Health Act 2010.

Results

Notifications and rates of pertussis in the 0–1 year age group were similar in both years, whereas notifications in older age-groups increased in 2015 compared to 2011 (Table 1). Conversely, the number of hospitalisations for children under 5 years was lower in 2015 compared to 2011 (Table 1).

Although source of illness data was missing or not known in at least 35% of cases for both years, during 2015 the most frequent source of illness for pertussis cases aged 0–4 years was another child at their childcare centre (Table 2). Siblings were the source of illness for approximately 20% of cases aged 0–4 years in both epidemic years (Table 2).

Case investigation consumed approximately 40 minutes of staff time per case, irrespective of the

Table 1. Comparison of pertussis notifications and hospitalisations in children aged 0–9 years in Sydney Local Health District, 2011 compared to 2015 (when enhanced surveillance was in place)

Period of follow-up and notification data	2011	2015*
Number of notifications by age group		
0–1yrs	43	49
2–4yrs	76	120
5–9yrs	172	248
Age-specific notification rate (per 100 000 population)		
0–1yrs	270	283
2–4yrs	371	522
5–9yrs	629	749
Number of hospitalisations by age group[†]		
0–1yrs	13	6
2–4yrs	2	1
5–9yrs	2	2
Age-specific hospitalisation rate (per 100 000 population)		
0–1yrs	81.7	34.7
2–4yrs	9.8	4.4
5–9yrs	7.3	6.4

* Notifications were extracted for the full calendar year of 2015. Rates were also calculated for the full 2015 calendar year.

† Hospitalisations with a respiratory-related cause, within 22 days of notification date.

NB: Numbers of notifications from NCIMS may change over time depending on the date of extraction from the database.

Source: Data from the NSW Communicable Disease Register, accessed through the Secure Analytics for Population Health Research and Intelligence (SAPHaRI) portal. Date of extraction and analysis 08/01/2018.

Table 2. Comparison of source of illness for pertussis cases aged 0–4 years in Sydney Local Health District in the year 2011 compared to 2015.

Source of illness for cases aged 0–4 years	2011		2015	
	N	%	N	%
Parent	9	7.6	9	5.2
Sibling	28	23.5	36	20.7
Other household (including grandparents)	4	3.4	2	1.1
Other non-household	12	10.0	20	11.5
Child at CCC	13	10.9	45	25.9
Unable to be identified/not stated	53	44.6	62	35.6
Totals	119	100	174	100

NB: Numbers of notifications in NCIMS may change over time depending on the date of extraction.

Source: Data from the Notifiable Conditions Incident Management System accessed through the Secure Analytics for Population Health Research and Intelligence (SAPHaRI) portal. Date of extraction and analysis: 18/01/2018

case's age or number of high-risk contacts. Of the 186 cases aged 5–9 years followed up during 2015, eight cases (4.3%) were identified with high-risk contacts requiring PEP (i.e. equated to a total 15.5 hours to identify one case with high-risk contacts), compared to routine follow-up of 129 cases aged 0–4 years, which identified 15 cases (11.7%) with high-risk contacts requiring PEP (5.7 hours to identify one case with high-risk contacts).

Discussion

The evaluation of our enhanced surveillance policy suggests that active follow-up of pertussis notifications in children aged 5–9 years during the epidemic year of 2015 may have led to lower hospitalisation rates for children aged 0–4 years when compared to the previous epidemic year of 2011, though notification rates of pertussis were higher in all age groups in 2015. Additionally, we identified a greater proportion of cases of pertussis in children aged 0–4 years likely acquired from another child at a childcare centre rather than a sibling, when compared to 2011. However, enhanced surveillance greatly increased PHU workload while identifying only a much smaller proportion of cases with high-risk contacts, compared to follow-up of cases

aged 0–4 years only. Therefore, our results suggest little incremental benefit from extending routine follow-up of children aged 0–4 years to those aged 5–9 years.

The lower hospitalisation rates for children aged 0–1 years during 2015 when compared to 2011 is noteworthy but confounded by the introduction of the maternal pertussis vaccination program in April 2015.¹² The significant protective effect of maternal vaccination for infants¹³ may have contributed to this lower rate. During 2015, the PHU also regularly sent out advice promoting testing to doctors and parents, which may have led to better detection and therefore higher notification rates for children overall.

Although siblings have been implicated as the most important source of pertussis for children aged 0–4 years⁷, our data indicates that during 2015, childcare centres were the most likely source of pertussis for these children. This epidemiological trend is unlikely to be sustained with new legislation requiring all children to be age-appropriately vaccinated prior to childcare enrolment³ and reintroduction of a pertussis booster for those aged 18 months on the national immunisation program.¹⁴

Our evaluation was limited by a small sample size and therefore lacking power to make confident statistical comparisons. Secondly, immunity from vaccination wanes over time¹ and changes in vaccination policy may have contributed to different levels of population susceptibility at different times, confounding any differences in data found in our evaluation. Thirdly, although we compared age cohorts of children offered the same vaccination schedule from birth, the comparison of data in 2011 versus 2015 was confounded by the introduction of the maternal pertussis vaccination program.¹²

Overall, our results suggest that there is little cost-benefit of actively following up pertussis cases aged 5–9 years, however, evaluation studies with larger cohorts of children over longer periods of time with more complete information on source of illness are needed to help confirm these results.

Author details

Kwendy Cavanagh¹

Emma Quinn¹

Alexandre S Stephens²

Zeina Najjar¹

Essi Huhtinen¹

Leena Gupta¹

1. Sydney Local Health District, Public Health Unit, Royal Prince Alfred Hospital, Missenden Road, Camperdown, Sydney, NSW, Australia

2. Northern NSW Local Health District, Research Office, Murwillumbah District Hospital, Ewing Street, Murwillumbah, NSW, Australia

Corresponding author

Dr Leena Gupta

Clinical Director, Public Health

Public Health Unit, Sydney Local Health District

Level 9 North, KGV Building, Missenden Road Camperdown, 2050

NSW Australia

Email: Leena.Gupta@health.nsw.gov.au

Tel: (+61) 02 9515 9420

Fax: (+61) 02 9515 9440

References

1. Misegades LK, Winter K, Harriman K, Talarico J, Messonnier NE, Clark TA, et al. Association of childhood pertussis with receipt of 5 doses of pertussis vaccine by time since last vaccine dose, California, 2010. *JAMA* 2012;308(20):2126–32.
2. Australian Technical Advisory Group on Immunisation (ATAGI). The Australian Immunisation Handbook 10th edition [Online]. Canberra: Australian Government Department of Health; 2015. Accessed on Jan 15 2017. Available from: <http://www.immunise.health.gov.au/internet/immunise/publishing.nsf/Content/Handbook10-home>
3. NSW Government. NSW Public Health Act, 2010, No.127 [Legislation]. Amended 2017. NSW Government: 2010. Accessed on May 1 2018. Available from: <https://www.legislation.nsw.gov.au/#/view/act/2010/127/fullh/view/act/2010/127/full>
4. NSW Health. Pertussis Control Guideline for Public Health Units [Guideline]. NSW Health 2015 Accessed Feb 25 2016. Available from: <http://www.health.nsw.gov.au/Infectious/controlguideline/Pages/pertussis.aspx>
5. Pillsbury A, Quinn HE, McIntyre PB. Australian vaccine preventable disease epidemiological review series: pertussis, 2006–2012. *Commun Dis Intell Q Rep* 2014;38(3):E179–194.
6. de Greeff SC, Mooi FR, Westerhof A, Verbaakel JMM, Peeters MF, Heuvelman CJ, et al. Pertussis disease burden in the household: how to protect young infants. *Clin Infect Dis* 2010;50(10):1339–45.
7. de Greeff SC, de Melker HE, Westerhof A,

Schellekens JFP, Mooi FR, van Boven M. Estimation of household transmission rates of pertussis and the effect of cocooning vaccination strategies on infant pertussis. *Epidemiology* 2012;23(6):852–60.

8. Jardine A, Conaty SJ, Lowbridge C, Thomas J, Staff M, Vally H. Who gives pertussis to infants? Source of infection for laboratory confirmed cases less than 12 months of age during an epidemic, Sydney, 2009. *Commun Dis Intell Q Rep* 2010;34(2):116–21.
9. Terry JB, Flatley CJ, van den Berg DJ, Morgan GG, Trent M, Turahui JA, et al. A field study of household attack rates and the effectiveness of macrolide antibiotics in reducing household transmission of pertussis. *Commun Dis Intell Q Rep* 2015;39(1):E27–33.
10. McCallum LK, Liu B, McIntyre P, Jorm LR. Estimating the burden of pertussis in young children on hospitals and emergency departments: a study using linked routinely collected data. *Epidemiol Infect* 2014;142(4):695–705.
11. National Health and Medical Research Council (NHMRC). National Statement on Ethical Conduct in Human Research (2007) (updated May 2015) [Guideline]. NHMRC 2015. Accessed 12 March 2018 Available from: <https://www.nhmrc.gov.au/book/national-statement-ethical-conduct-human-research>
12. Beard F. Pertussis immunisation in pregnancy: a summary of funded Australian state and territory programs. *Commun Dis Intell Q Rep* 2015;39(3):E329–336
13. Amirthalingam G, Andrews N, Campbell H, Ribeiro S, Kara E, Donegan K, et al. Effectiveness of maternal pertussis vaccination in England: an observational study. *Lancet* 2014;384(9953):1521–8.
14. Australian Government Department of Health. National Immunisation Program

Schedule [last updated 08/12/2017]. Australian Government: 2017. Accessed on 01/05/2018. Available from: <https://beta.health.gov.au/topics/immunisation/immunisation-throughout-life/national-immunisation-program-schedule>